

Patient Information for Consent

OS13 Wrist Fracture Surgery

Expires end of July 2021

Mr. Z.O.Rahimtoola, MD, PhD, European Diploma Hand Surgery

Consultant Orthopaedic Hand Surgeon

www.GlobalHandSurgery.com

Get more information and references at www.aboutmyhealth.org

Tell us how useful you found this document at www.patientfeedback.org



The Information Standard  Certified Member

eidohealthcare.com

EIDO
HEALTHCARE

UNITED KINGDOM

Information about COVID-19 (Coronavirus)

On 11 March 2020 the World Health Organization confirmed COVID-19 (coronavirus) has now spread all over the world (this means it is a 'pandemic'). Even though lockdown has been eased, there is still a risk of catching coronavirus. Hospitals have very robust infection control procedures, however, it is impossible to make sure you don't catch coronavirus either before you come into the hospital or once you are there. You will need to think carefully about the risks associated with the procedure, the risk of catching coronavirus while you are in hospital, and of not going ahead with the procedure at all. Your healthcare team can help you understand the balance of these risks. If you catch the coronavirus, this could affect your recovery and might increase your risk of pneumonia and even death. Talk to your healthcare team about the balance of risk between waiting until the pandemic is over (this could be many months) and going ahead with your procedure.

Please visit the World Health Organization website: <https://www.who.int/> for up-to-date information.

Information about your procedure

It is essential that your surgery goes ahead as soon as possible. Following the Covid-19 (coronavirus) pandemic, some of the hospital processes have changed. You will need a coronavirus test before your operation. This involves the healthcare team taking a nasal and throat swab (using cotton wool to take a sample from the surface of your nasal passage and throat). The operation may need to go ahead before the results are available. Your healthcare team can tell you about the risks of coronavirus.

Coronavirus is highly contagious (meaning it spreads easily from person to person). The most common way that people catch it is by touching their face after they have touched a person or surface that has the virus on it. Try not to touch your face, especially if you have not washed your hands.

If possible, wash your hands with alcoholic gel or soap and water when you enter the hospital, at regular intervals after that, and when you move from one part of the hospital to another.

Be aware of social distancing. Chairs and beds are spaced apart. If your healthcare team need to be close to you, they will wear personal protective equipment (PPE). If you can't hear what they are saying because of their PPE, ask them to repeat it until you can.

You may not be allowed visitors, or your visiting may be restricted.

Your surgery is essential and the hospital and health professionals looking after you are very well equipped to perform it in a safe and clean environment. Guidance about coronavirus may change quickly; your healthcare team will have the most up-to-date information.

What is a wrist fracture?

A wrist fracture is a break of one or both of the bones in your forearm near your wrist joint. A wrist fracture is sometimes known as a 'distal radius fracture'.

Your surgeon has recommended an operation to treat your broken wrist. However, it is your decision to go ahead with the operation or not.

This document will give you information about the benefits and risks to help you to make an informed decision. If you have any questions that this document does not answer, ask your surgeon or the healthcare team.

How does a wrist fracture happen?

Many wrist fractures happen to older people when they fall with their hand stretched out. Wrist fractures can also happen to younger people.

Sometimes the fracture is just a little crack in the bone. However, the fracture can be more severe with the bone broken in many places and damage to the surface of your wrist joint.

What are the benefits of surgery?

The aim is to hold the pieces of bone in a good position while the fracture heals. This should help your wrist to work better.

Are there any alternatives to surgery?



Some wrist fractures heal well in a cast. Before placing your wrist in a cast, you may need to have the bones pulled into a better position (called a manipulation). You will need a local or general anaesthetic.

If your wrist is badly broken, the bones do not always stay in a good position in the cast. Your surgeon may recommend an operation.

What will happen if I decide not to have the operation?

You may need to have a manipulation and then your wrist will be put in a cast. If the bones do not heal in a good enough position, you are more likely to have problems with your wrist in the future.

It is usually only possible to perform the operation within about 2 weeks of the injury.



- a Wires
- b Plate and screws
- c An external fixator

What does the operation involve?

The healthcare team will carry out a number of checks to make sure you have the operation you came in for and on the correct side. You can help by confirming to your surgeon and the healthcare team your name and the operation you are having.

Various anaesthetic techniques are possible. Your anaesthetist will discuss the options with you.

You may also have injections of local anaesthetic to help with the pain after the operation. You may be given antibiotics during the operation to reduce the risk of infection.

There are several different ways of fixing a broken wrist.

- Wires inserted through your skin.
- Plates fixed to the bone with screws.
- An 'external fixator' (using a frame and pins).

Your surgeon will discuss which option, or combination of options, is best for you.

They will close your skin with stitches or clips and place a bandage or cast on your wrist.

What should I do about my medication?

Make sure your healthcare team knows about all the medication you take and follow their advice. This includes all blood-thinning medication as well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter. Anti-inflammatory painkillers may prevent the fracture from healing properly, so it is better not to take these if possible.

What can I do to help make the operation a success?

If you smoke, stopping smoking may reduce your risk of developing complications and will improve your long-term health. Nicotine is known to stop fractures from healing.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

You can reduce your risk of infection in a surgical wound.

- Try to have a bath or shower either the day before or on the day of the operation. If you have a temporary cast, you must keep it dry.
- Keep warm around the time of the operation. Let the healthcare team know if you feel cold.

What complications can happen?

The healthcare team will try to reduce the risk of complications.

Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you.

Some complications can be serious.

You should ask your doctor if there is anything you do not understand.

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

General complications of any operation

- Pain. The healthcare team will give you medication to control the pain and it is important that you take it as you are told so you can move about as advised.
- Bleeding during or after the operation.
- Infection of the surgical site (wound). It is usually safe to shower after 2 days but you should check with the healthcare team. Keep your wound dry and covered. If you have a cast, you must keep it dry. Let the healthcare team know if you get a high temperature, notice pus in your wound, or if your wound becomes red, sore or painful. An infection usually settles with antibiotics but you may need another operation.
- Unsightly scarring of your skin.
- Difficulty passing urine. You may need a catheter (tube) in your bladder for 1 to 2 days.

Specific complications of this operation

- Damage to nerves, leading to a patch of numb skin or a tender scar (risk: 1 in 10). This usually gets better but may be permanent.
- Infection around a wire (risk: 1 in 10). This usually settles when the wire is removed. You may need a course of antibiotics.
- Infection of the bone caused by an infection around a wire or plate (risk: 1 in 250). You will need a course of antibiotics or sometimes another operation.
- The bones move out of position. You may need another operation to fix the bones in a good position again.
- Severe pain, stiffness and loss of use of your wrist and hand (complex regional pain syndrome) (risk: 1 in 10). The cause is not known. You may need further treatment including painkillers and physiotherapy. Your wrist and hand can take months or years to improve. You may be able to reduce this risk by taking a 500mg vitamin C tablet each day for 6 weeks after the operation.
- Tendon problems affecting your thumb. This can cause pain when you move your thumb. One of the tendons that moves your thumb can sometimes snap after a few weeks. You will need an operation to redirect another tendon so you can move your thumb.

- Carpal tunnel syndrome, where there is increased pressure on the nerve that crosses the front of your wrist (the median nerve) (risk: 1 in 20). This results in pain or numbness in your thumb, index and middle fingers. You may need a carpal tunnel release operation to relieve the pressure.

How soon will I recover?

In hospital

After the operation you will be transferred to the recovery area and then to the ward.

Keep your wound dry for 4 to 5 days, and use a waterproof dressing when you have a bath or shower.

The healthcare team will tell you if you need to have any stitches removed or dressings changed.

You should be able to go home after 1 to 2 days. However, your doctor may recommend that you stay a little longer.

If you are worried about anything, in hospital or at home, contact the healthcare team. They should be able to reassure you or identify and treat any complications.

Returning to normal activities

The healthcare team will tell you when you can return to normal activities.

Keep your hand raised for the first week so that the swelling settles. It is important to move your fingers to help reduce any stiffness.

The fracture usually heals in about a month. If your surgeon used wires or an external fixator, they will usually remove these in the outpatient clinic. This procedure does not need another anaesthetic.

You may need some physiotherapy if your wrist is stiff.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Do not drive until you are confident about controlling your vehicle and always check your insurance policy and with your doctor.

The future

Most people make a good recovery and are able to return to normal activities. However, recovery can take many months.

Your doctor may recommend tests or further treatment to reduce the risk of another fracture.

- Looking into any cause for your fall, such as a dizzy spell or blackout.
- An exercise programme to improve your balance and muscle strength.
- Medication to make your bones stronger if you have osteoporosis (brittle bones).

The injury can cause you to lose some wrist movement permanently and sometimes cause your grip to be weaker. About 1 in 25 people develops arthritis in their wrist but this does not often need any treatment.

Summary

For some types of wrist fracture, an operation is the best way to make sure the bones heal in a good position.

Surgery is usually safe and effective but complications can happen. You need to know about them to help you to make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Acknowledgements

Reviewers: Stephen Milner (DM, FRCS), Adam Watts (MBBS, FRCS)

Illustrator: Medical Illustration Copyright © Medical-Artist.com

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.